



FIBROIDS & FIBROID TREATMENT OPTIONS

CHANEL, 30
Procedure Date
April 2019

PATRICE, 49
Procedure Date
June 2018

RHONDA, 36
Procedure Date
December 2017

Fibroids 101

Fibroids 101

What are fibroids?


Fibroids are noncancerous tumors that grow in the smooth muscle cells of the uterus.

Where do fibroids come from?¹

The cause of fibroids isn't well understood. Risk factors include but are not limited to age, race, obesity, and family history of uterine fibroids.

Are fibroids common?^{2,3}

Approximately 3 million women are diagnosed with fibroids in the US.



Symptoms⁴

- Heavy and painful or prolonged periods
- Stomach, lower-back and pelvic pain
- Anemia
- Stomach protrusion (causing a woman to look pregnant when she is not)
- G.I. issues like gas and constipation
- Infertility
- Painful sex
- Frequent urination


80% of African American women develop fibroids by the age of 50.⁵

3.6 Women with fibroids waited an average of 3.6 years to seek treatment.⁶
YEARS

3M 3 million women^{2,3}
3 Million women are diagnosed with symptomatic uterine fibroids in the US.

Types of Fibroids

BASED ON LOCATION



Fibroids are classified by their location in the uterus:

Subserosal*

Outer wall of the uterus and typically cause bulk or pressure symptoms.

Intramural*

Within the muscular walls of the uterus and typically can cause heavy bleeding or pressure symptoms.

Submucosal*

Either inside or abutting the uterine cavity and typically cause heavy bleeding (least common type).

Pedunculated

Fibroids on a thin stalk (less common)

***Fibroid types that can be treated with the Acessa procedure (also underlined on graphic).**

Fibroid Size

APPROXIMATE FIBROID SIZE COMPARISON CHART

Fibroids may range in size from a pea to a melon.⁷ Typically patients are provided two measurements.

Fibroid size – Often expressed as centimeters (cm) by diameter.

Uterus size – Often expressed as the number of weeks pregnant the woman would be if the uterus + fibroids was the size of the uterus + fetus. This chart does not take into account the presence of multiple fibroids. Multiple fibroids can cause the uterus size to be larger than in the case of a single fibroid.

Symptoms and severity depend on both the location and size of the fibroid(s)⁸

 Sweet Pea/Apple Seed UTERUS SIZE: 6 weeks DIAMETER: 0.5 cm	 Blueberry UTERUS SIZE: 7 weeks DIAMETER: 1 cm	 Raspberry/Cranberry UTERUS SIZE: 8 weeks DIAMETER: 2 cm	 Strawberry UTERUS SIZE: 10 weeks DIAMETER: 3 cm
 Lime UTERUS SIZE: 11 weeks DIAMETER: 5 cm	 Peach UTERUS SIZE: 13 weeks DIAMETER: 8 cm	 Lemon UTERUS SIZE: 14 weeks DIAMETER: 9 cm	 Mango UTERUS SIZE: 19 weeks DIAMETER: 15 cm
 Spaghetti Squash UTERUS SIZE: 21 weeks DIAMETER: 27 cm	 Lettuce UTERUS SIZE: 27 weeks DIAMETER: 35 cm	 Honeydew UTERUS SIZE: 32 weeks DIAMETER: 42 cm	 Watermelon UTERUS SIZE: 36 weeks DIAMETER: 50 cm

This table is used to illustrate approximate sizes and diameters of a single fibroid based on average fruit sizes. It does not reflect clinical data. Source: Physician Working Group. The safety and effectiveness of the Acessa procedure has not been evaluated in women with uterine size > 14 weeks.

Other Conditions

THAT ARE COMMONLY FOUND WITH FIBROIDS

Endometriosis⁹

A condition resulting from the appearance of endometrial tissue in sites outside the uterine cavity, such as the ovaries or pelvic wall.

Adenomyosis¹⁰

A type of endometriosis. Presence of endometrial tissue inside the muscle layers of the uterus. May be focal or diffuse. Focal creates focal adenomyomas; often softer than a fibroid with a poorly defined border. Diffuse results in a uniformly enlarged uterus.


Polyps¹¹

A benign overgrowth of endometrial tissue from the uterine lining. They attach to the uterine lining by a large base or thin stalk.

Cysts


A closed, sac-like structure, typically filled with liquid, semisolid or gaseous material. May be benign or malignant and typically forms on the ovaries.

The Acessa Procedure is indicated for the treatment of symptomatic uterine fibroids. It should not be used to treat any of the conditions listed above. Talk to your doctor to determine the right treatment option(s) for you.




Fibroid Treatment Options


Fibroid Treatment Options



Hysterectomy and Myomectomy can be performed using different approaches




The Acesa procedure incision points



This chart is representative only, to be used in shared decision making between patients and physicians. It is based on physician and patient input. The invasiveness and time to return to daily life of different procedures is dependent on numerous factors. This chart is not intended to be a representation of a single study or clinical data.


Fibroid Treatment Options




This chart is representative only, to be used in shared decision making between patients and physicians. It is based on physician and patient input. The invasiveness and time to return to daily life of different procedures is dependent on numerous factors. This chart is not intended to be a representation of a single study or clinical data.

Hysteroscopic Procedures

Hysteroscopic Myomectomy¹²




- > Removal of fibroids from uterus through the vagina using a scope (camera) to visualize the fibroid and a tool to remove the fibroid.
- > Also known as hysteroscopy or hysteroscopic resection.
- > Addresses fibroids that are in the uterine cavity; these fibroids typically cause heavy bleeding.
- > Typically a short recovery time.



Endometrial Ablation¹³


DESIGNED FOR ABNORMAL UTERINE BLEEDING



- > Designed to address abnormal uterine bleeding.
- > Destroys the uterine lining (endometrium) using heated fluids, microwave energy, extreme cold, or high-energy radiofrequencies.
- > Also known as GEA (global endometrial ablation).
- > Can be effective for multiple causes of abnormal uterine bleeding.


The Acesa Procedure

Lap-RFA^{14,15,16}




Prep

You'll be brought into the operating room for anesthesia




Access

After you're asleep, your physician will make a minimum of three small incisions for a camera, ultrasound, and Acesa handpiece.




Visualize

Your physician uses a special ultrasound directly on the uterus that allows them to find and ultimately treat more fibroids than other standard imaging methods.



Deploy

Next your physician deploys the tip of the Acesa handpiece into the fibroid and only the fibroid.



Treat

Controlled heat will destroy fibroid tissue. The heat shrinks the tissue mass into a fibroid into a soft mass that is absorbed by the body over time. Your physician will repeat this process until every targeted fibroid is treated.



Recover

After surgery, most patients get cleared to go home in two hours, and return to work in 4-5 days. Most women experience relief in the first 3 months, and continual improvement for 12 months.






Lap-RFA

How Acessa Works

The Acessa procedure is performed laparoscopically through small incisions in the abdomen. The procedure has an ultrasound probe and allows for the use of a camera, which helps physicians visualize fibroids.

The heat causes fibroid tissue to shrink overtime, resulting in symptom improvement. Women typically see the most improvement within 3-6 months with continued improvement throughout the first year.¹⁵ Different symptoms resolve at different rates; some may resolve faster or slower than others.¹⁵

Coagulative Necrosis → Destroys Fibroid Tissue




White colored fibroids are representative of treated fibroid tissue

Lap-RFA


Radiofrequency ablation (RFA) is not new

Other procedures that treat conditions in the liver, bone and prostate use the same technology, but the Acessa procedure is the first of its kind to use **Lap-RFA** on uterine fibroids. Although the risk is minimal, there is still a chance of skin burns.

Other rare but serious risks of the Acessa procedure include, but are not limited to, infection, internal injury, blood loss and complications related to laparoscopic surgery and/or general anesthesia.



Acessa Procedure equipment setup in the Operating Room



Results may vary.

VOLUMETRIC SHRINKAGE EXAMPLE


Acessa is designed to shrink the volume of a fibroid.

Even a seemingly small decrease in diameter yields a large decrease in volume.

- > Clinical data showed an average **45% reduction** in fibroid volume over 12 months, although results may vary.¹⁵
- > Volume = $\frac{4}{3} \pi r^3$
- > Acessa destroys fibroid tissue and causes shrinkage over time, which helps relieve symptoms such as heavy periods, pelvic pressure, urinary frequency, etc.¹⁵

Other volumetric shrinkage examples.


This image shows the relationship between diameter reduction size and volume reduction size for an initial fibroid diameter of 5 cm.



Results may vary and patients may still experience symptoms associated with their fibroids despite a reduction in volume.

UAE




Uterine Artery Embolization (UAE)



- > Interventional Radiologist uses embolizing agent to stop blood flow to fibroids.
- > Can be performed under conscious sedation (while awake).
- > Common side effects may include: post-embolization syndrome, amenorrhea, pain, discharge, angiography-related complications, vaginal discharge of fibroid material, hot flashes, endometritis/myometritis, deep leg vein thrombosis/pulmonary embolus.¹⁷
- > Works via ischemic necrosis versus coagulative necrosis (LAP RFA).

Myomectomy

Myomectomy



- > Opening of the uterus to remove fibroids by excising them from the uterus.
- > Depending on the size, location, and number of fibroids, can be performed:
 - Open (abdominal incision)
 - Laparoscopic (Robotic or straight stick)*
- > Typically a 2-4 week recovery for laparoscopic and abdominal hysterectomy.¹⁸






Illustration of hysteroscopic myomectomy.
Refer to page 11

Hysterectomy

Hysterectomy



- > Removal of the uterus.
- > Depending on the size, location, and number of fibroids, can be performed:
 - Open (abdominal incision)
 - Laparoscopic (Robotic or straight stick)*
 - Vaginal*
- > Typically a 3-6 week recovery.¹⁹

*Often these options will be referred to as a “minimally invasive” hysterectomy

Disclaimer

This booklet has described several different options for the treatment of symptomatic fibroids. Not every procedure may be right for you. You should discuss your treatment options, as well as the risks and benefits associated with each treatment option, with your physician.

Only you and your physician can determine the right course of treatment for you.

REFERENCES

1. Johns Hopkins – Fibroids <https://www.hopkinsmedicine.org/health/conditions-and-diseases/uterine-fibroids>
2. Wise LA, Laughlin-Tommaso SK. Clin Obstet Gynecol. 2016;59(1):2-24
3. Alexander AL, et al. Obstet Gynecol. 2019;133(1):6-12
4. Mayo Clinic – <https://www.mayoclinic.org/diseases-conditions/uterine-fibroids/symptoms-causes/syc-20354288>
5. Marsh, E., Al-Hendy, A., Kappas, D., Galitsky, A., Stewart, E. A., & Kerolous, M. (2018). Burden, Prevalence, and Treatment of Uterine Fibroids: A Survey of U.S. Women. Journal of women's health (2002), 27(11), 1359–1367. <https://doi.org/10.1089/jwh.2018.7076>
6. Borah, Bijan J. et al. The impact of uterine leiomyomas: a national survey of affected women, American Journal of Obstetrics & Gynecology, October 2013, Volume 209, Issue 4, 319.e1–319.e20. Wise LA, Laughlin-Tommaso SK. Clin Obstet Gynecol. 2016;59(1):2-24. Alexander AL, et al. Obstet Gynecol. 2019;133(1):6-12
7. Johns Hopkins Medicine. (2017, May 22). Uterine Fibroids. Retrieved from https://www.hopkinsmedicine.org/gynecology_obstetrics/specialty_areas/gynecological_services/conditions/fibroids.html
8. UNC School of Medicine. (2020). Fibroid Tumors. Retrieved from <https://www.med.unc.edu/obgyn/migs/our-services/what-kinds-of-problems-do-the-doctors-in-this-group-treat/fibroid-tumors/>
9. Mayo Clinic – Endometriosis <https://www.mayoclinic.org/diseases-conditions/endometriosis/symptoms-causes/syc-20354656>
10. Diffuse and Focal Adenomyosis: MR Imaging Findings Jae Young Byun, Sung Eun Kim, Byung Gil Choi, Gi Young Ko, Seung Eun Jung, and Kyu Ho Choi RadioGraphics 1999 19:suppl_1, S161-S170 https://pubs.rsna.org/action/showCitFormats?doi=10.1148%2Fradiographics.19.suppl_1.g99oc03s161
11. Mayo Clinic - Uterine Polyps https://www.mayoclinic.org/diseases-conditions/uterine-polyps/symptoms-causes/syc-20378709_Z, Firtina Tuncer, S., Akpınar, F., Kayıkçıoğlu, F., & Koç, S. (2016). Prevalence of endometrial polyps coexisting with uterine fibroids and associated factors. Turkish journal of obstetrics and gynecology, 13(1), 31–36. <https://doi.org/10.4274/tjod.36043>
12. Myomectomy. (2019, June 11). Retrieved from <https://www.mayoclinic.org/tests-procedures/myomectomy/about/pac-20384710>
13. Endometrial ablation. (2018, September 8). Retrieved from <https://www.mayoclinic.org/tests-procedures/endometrial-ablation/about/pac-20393932>
14. Acesa Health, Acesa ProVu System Instructions for Use, Nov. 2018
15. Chudnoff SG, Berman J, Levine DJ, Harris M, Guido RS, Banks E. SG Chudnoff, et al. Outpatient Procedure for the Treatment and Relief of Symptomatic Uterine Myomas. Obstetrics and Gynecology, 2013;121(5):1075–82. Median of 5 missed work days.
16. Galen DI, Pemueler RR, Leal JG, Abbott KR, Falls JL, Macer J. Laparoscopic radiofrequency fibroid ablation: phase II and phase III results. JLS. 2014;18(2):182-190. doi:10.4293/108680813X13693422518353. Median of 4 missed work days.
17. Kröncke, T., David, M., & participants of the Consensus Meeting (2015). Uterine Artery Embolization (UAE) for Fibroid Treatment – Results of the 5th Radiological Gynecological Expert Meeting. Geburtshilfe und Frauenheilkunde, 75(5), 439–441. <https://doi.org/10.1055/s-0035-1545999>
18. Laughlin-Tommaso SK, Lu D, Thomas L, Diamond MP, Wallace K, Wegienka G, Vines AI, Anchan RM, Wang T, Maxwell GL, Jacoby V, Marsh EE, Spies JB, Nicholson WK, Stewart EA, and Myers ER. Short-term quality of life after myomectomy for uterine fibroids from the COMPARE-UF Fibroid Registry. Am J Obstet Gynecol 2020; 222(4):345.e341-345.e322
19. Consultant, C.O., Lingman, G. and Ottosen, L. (2000). Three methods for hysterectomy: a randomised, prospective study of shortterm outcome. BJOG: An International Journal of Obstetrics & Gynaecology, 107: 1380-1385. <https://doi.org/10.1111/j.1471-0528.2000.tb11652.x>

Important Safety Information The Acesa ProVu system is intended to identify and shrink symptomatic uterine fibroids. The Acesa ProVu system is used by trained physicians during laparoscopic surgery under general anesthesia. Rare but serious risks of this procedure include, but are not limited to, infection, internal injury, blood loss and complications related to laparoscopic surgery and/or general anesthesia. This procedure is not recommended for women who are planning future pregnancy. This information is not medical advice. Please discuss the risks and benefits with your doctor to find out if the Acesa procedure may be right for you.

PP-02217 Rev 002 ©2020 Hologic, Inc. All Rights Reserved. Hologic, Acesa, Acesa ProVu, MyoSure, The Science of Sure and associated logos are trademarks and/or registered trademarks of Hologic, Inc. and/or its subsidiaries in the United States and/or other countries.



HOLOGIC®  **acessa®**

gynsurgicalsolutions.com/patients/treatment-options/acessa/procedure-steps-and-components/

SHARON, 41
Procedure Date
November 2019